Hazard Evaluation for the Mars Pathfinder Prime Landing Site from Synthesis of Radar Ranging and Continuous Wave Observations during the -1995 Mars Opposition

M. A. Slade, R. F. Jurgens (JPL/Caltech), A. F. C. Haldemann (Caltech), and D. L. Mitchell (JPL/Caltech)

During the 1995 Mars Opposition, the sub-Earth latitudes available for Earthbased radar observations fortuitously covered a large percentage of the possible landing sites for Mars Pathfinder. The preliminary prime landing site at the mouth of Ares Vallis in southeastern Chryse Planitia in particular was available at closest approach when the radar returns would be strongest. The 100 x 200 km Ares Vallis landing ellipse covers an area that was rejected as a Ving Lander site in 1976 due to Goldstone CW radar echoes" with low signal-to-noise (SNR). The Goldstone radar system at 3.5-cm is now 12 dB more sensitive than the 13-cm system in use at the time of the Viing Ianding site assessment, and thus definitive results on the potential problems of this site are expected. We will present updated and integrated evaluations for both landing hazards and Rover operation at the Ares Vallis site based on two types of radar observations: radar delay-Doppler profiles (ranging for short) and Continuous Wave (CW) radar spectra obtained using the Goldstone 70-m antenna. at X-band (3.5-cm). The "ranging" data are in the form of delay-Doppler profiles which fall along a "great circle" on Mars which are nearly lines of constant latitude. The Doppler frequency resolution provides a longitude resolution of about 4 km, while the delay resolution implies a latitude resolution of about 100 km. Three quantities can be extracted from the multiple views of the pieces of the profiles: topography, reflectivity, and Hagfor's C parameter (a measure of rms slope). For CW observations, echoes are recorded simultaneously in both the same sense of circular polarization as transmitted (the SC sense) and the opposite (OC) sense. The circular polarization ratio thus obtained is a measure of near-surface complexity, or "roughness," at scales near the 3.5-cm observing wavelength.

DPS Category 6 Running # Session 0.00 Oral preferred Poster preferred Either 🔲 1s this your first DPS presentation? Yes No Would you be willing to act as Session Chair? Yes X Is your abstract newsworthy, and if so, would you be willing to prepare a news release and be available for interviews with reporters? Yes Maybe X Paper presented by Martin A. Slade Jet Propulsion Laboratory Mail Stop 238-420 4800 Oak Grove Drive Pasadena, CA 91109-8099 Phone: 818-354-2765 Fax: 818-354-6825 Email: marty@bvd.jpl.nasa.gov Special instructions: Membership Status (First Author): DPS-AAS Member Non-Member Student Member Student Non-Member Sponsor:

Division for Planetary Sciences Abstract Form

Abstract submitted for DPS [Division for Planetary Sciences] meeting

Date submitted: LPI electronic form version 5/95